



Grosafe
garden



EnSpray 99[®]

SPRAYING OIL

INSECTICIDE, MITICIDE & FUNGICIDE

An emulsifiable, narrow range spray oil for the control of scale, aphids and mites on citrus, pip and stone fruits, thrips on citrus, European red mite, mealy bug, scale insects and aphids on ornamentals and roses. Controls powdery mildew on grapes.

SCE5482/F09

200ml

makes
up to
20L

CONTROLS



Aphids



European
red mite



Mealy bugs



Mites



Powdery
Mildew



Scale



Thrips

See back label for full list

EnSpray 99 200ml

PEEL HERE



DANGER KEEP OUT OF REACH OF CHILDREN READ LABEL BEFORE USE
HAZARD CLASSIFICATIONS: Aspiration hazard Category 1,
Eye irritation Category 2, Designed for biocidal action.

GENERAL INFORMATION: ENSPRAY 99 is a highly refined narrow distillation range spraying oil which contains emulsifiers to facilitate mixing with water.

DIRECTIONS FOR HOME GARDEN USE

Mixing Instructions: Ensure the sprayer is clean. Pour the required amount of water into the sprayer then add ENSPRAY 99 and agitate well. When mixing with other pesticides add the other product first, according to the manufacturer's instructions, then add the ENSPRAY 99 last. Maintain continuous agitation while spraying. The rates suggested are for high volume spraying to the point of run-off.

RECOMMENDATIONS FOR USE:

| PLANTS | USE | RATE |
|--|---|--|
| Deciduous plants: Apples, pears and stone fruit | For control of European red mite, San Jose scale and aphids | 15-20ml per litre of water. For scale control apply during dormant period up until green tip |
| Citrus | For the control of scale insects, aphids, citrus red mite and thrips | 10ml per litre of water |
| Evergreen Plants: Roses and Ornamentals | For the control of European red mite, mealy bug, scale insects and aphids | |
| Grapes | Powdery mildew control | 20ml per litre of water. Repeat at 7-10 day intervals. |
| If you have any doubt about crop safety contact Sipcam New Zealand Ltd, or apply to a small area first. | | |
| Indoor plants | Leaf polish. Apply with a soft cloth | 5ml per 200ml (1 cup) of water. |

WITHHOLDING PERIOD: 1 day

Registered to and distributed by
SIPCAM NEW ZEALAND LTD
20 Jean Batten Drive, Mt Maunganui, 3116
PO Box 14 - 450 Tauranga Central 3141
Tel: 0800 220002
Email: info@sipcam.co.nz
www.grosafegarden.co.nz

See container for Batch code
& Date of Manufacture



9 421902 986482 >

TOXICITY: DANGER - May be fatal if swallowed and enters airways. Serious eye irritant.

ECOTOXIC: Designed for biocidal action. Do not apply directly into or onto water.

Avoid contamination of any water supply with product or empty container.

PERSONAL PROTECTION: When mixing or loading wear overalls, impervious gloves and eye/ face protection. Avoid breathing spray mist. Wash hands and exposed skin after work.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from foodstuffs.

SPILLS AND DISPOSAL: When dealing with spills wear gloves. Contain and absorb with inert absorbent material then collect into container for disposal.

PRODUCT DISPOSAL: If possible dispose of by using according to the label, otherwise dispose of in an approved landfill.

CONTAINER DISPOSAL: Triple rinse container and add residue to mixture. Dispose of via an approved recycler or in rubbish collection.

SAFETY DATA SHEET: A safety data sheet is available from Sipcam New Zealand Ltd.

FIRST AID: For advice call a doctor or the National Poisons Centre (0800 764 766). If medical advice is needed, have product container or label at hand. If swallowed, DO NOT induce vomiting. Immediately call the National Poisons Centre or doctor for advice. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention. If on skin, wash with plenty of soap and water. If irritation occurs, get medical advice.

EMERGENCY RESPONSE 0800 CHEMCALL: (0800 243 622)

It is an offence to use this product on animals.

Restraints

DO NOT spray under conditions that cause slow drying: i.e. low temperatures, high humidity. DO NOT spray in the late evening or early morning or during the hours of darkness: this could cause damage to the fruit.

DO NOT use additional spray adjuvants.

COMPATIBILITY

DO NOT apply **ENSPRAY 99** within 14 days of applications of sulphur or lime sulphur, once crop growth has commenced.

Consult the label of the proposed additive. If doubt remains DO NOT mix products.

ACTIVE INGREDIENT: 821g/L (980 ml/L) mineral oil in the form of an emulsifiable concentrate. Unsulphonated residue not less than 99.0%.

Registered pursuant to the ACVM Act 1997, No P009104.
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to HSNO Act 1996, No HSR 100946.
See www.epa.govt.nz for approval conditions.

NOTICE TO BUYER: This product must be used in accordance with the directions for use. Other than warranted by New Zealand consumer protection law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.